

Existence of positive solutions for fractional q -difference equations involving integral boundary conditions with p -Laplacian operator

Yizhu Wang¹, Chengmin Hou^{1*}

(Department of Mathematics, Yanbian University, Yanji, 133002, P.R. China)

ABSTRACT: The existence of positive solutions is considered for a fractional q -difference equation with p -Laplacian operator in this article. By employing the Avery-Henderson fixed point theorem, a new result is obtained for the boundary value problems.

KEY WORDS: positive solutions; Riemann-Liouville fractional q -derivatives; Caputo fractional q -derivatives; p -Laplacian; Avery-Henderson fixed point theorem

I. INTRODUCTION

Fractional calculus is the extension of integer order calculus to arbitrary order calculus. With the development of fractional calculus, fractional differential equations have wide applications in the modeling of different physical and natural science fields, such as fluid mechanics, chemistry, control system, heat conduction, etc. There are many papers concerning fractional differential equations with the p -Laplacian operator [1-5] and fractional differential equations with integral boundary conditions [6-10].

In [11], Yang studied the following fractional q -difference boundary value problem with p -Laplacian operator

$$\begin{aligned} {}^c D_q^\beta (\phi_p ({}^c D_q^\alpha x(t))) &= f(t, u(t)), 0 < t < 1, 2 < \alpha < 3, \\ x(0) = x(1) &= 0, \quad {}^c D_q^\alpha x(1) = {}^c D_q^\alpha x(0) = 0, \end{aligned}$$

where $1 < \alpha, \beta \leq 2$. The existence results for the above boundary value problem were obtained by using the upper and lower solutions method associates with the Schauder fixed point theorem.

In [12], Yuan and Yang considered a class of four-point boundary value problems of fractional q -difference equations with p -Laplacian operator

$$\begin{aligned} D_q^\beta (\phi_p (D_q^\alpha x(t))) &= f(t, u(t)), 0 < t < 1, 2 < \alpha < 3, \\ x(0) = 0, x(1) &= ax\xi, \quad D_q^\alpha x(0) = 0, D_q^\alpha x(1) = bD_q^\alpha x(\eta), \end{aligned}$$

Where D_q^β, D_q^α are the fractional q -derivative of the Riemann-Liouville type with $1 < \alpha, \beta \leq 2$. By applying the upper and lower solutions method associated with the Schauder fixed point theorem, the existence results of at least one positive solution for the above fractional q -difference boundary value problem with p -Laplacian operator are established.

Motivated by the aforementioned work, this work discusses the existence of positive solutions for this fractional q -difference equation:

$$\begin{cases} D_{q,0^+}^\beta [\phi_p({}^c D_{q,0^+}^\alpha u(t))] + f(t, u(t)) = 0, t \in (0,1), \\ \phi_p({}^c D_{q,0^+}^\alpha u(0)) = [\phi_p({}^c D_{q,0^+}^\alpha u(0))] = \phi_p({}^c D_{q,0^+}^\alpha u(1)) = 0, \\ D_{q,0^+}^2 u(0) = D_{q,0^+} u(1) = 0, \\ au(0) + bD_{q,0^+} u(0) = \int_0^1 g(t)u(t)d_q t, \end{cases} \quad (1.1)$$

where $2 < \alpha \leq 3$, $2 < \beta \leq 3$, and $5 < \alpha + \beta \leq 6$. $\phi_p(u) = |u|^{p-2}u$, $p > 1$. ${}^c D_{q,0^+}^\alpha$ is the Caputo fractional q -derivative, $D_{q,0^+}^\beta$ is the Riemann-Liouville fractional q -derivative.

We will always suppose the following conditions are satisfied:

(H₁) $g(t) : [0,1] \rightarrow [0, +\infty)$ with $g(t) \in L^1[0,1]$, $\int_0^1 g(t)d_q t > 0$, and $\int_0^1 tg(t)d_q t > 0$;

(H₂) $a, b \in (0, +\infty)$, $a > \int_0^1 g(t)d_q t$ and $b > a$;

(H₃) $f(t, u) : [0,1] \times (0, +\infty) \rightarrow (0, +\infty)$ is continuous.

II. BACKGROUND AND DEFINITIONS

To show the main result of this work, we give in the following some basic definitions and theorems, which can be found in [13,14].

Definition 2.1 [13] Let $\alpha \geq 0$ and f be a function defined on $[0,1]$. The fractional q -integral of Riemann-Liouville type is $(I_{q,0^+}^0 f)(t) = f(t)$ and

$$(I_{q,0^+}^\alpha f)(t) = \int_0^t \frac{(t-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s)d_q s, \quad \alpha > 0, t \in [0,1].$$

Lemma 2.2 [13] Let $\alpha > 0$, then the following equality holds:

$$I_{q,0^+}^\alpha {}^c D_{q,0^+}^\alpha f(t) = f(t) - \sum_{k=0}^{[\alpha]-1} \frac{t^k}{\Gamma_q(k+1)} (D_{q,0^+}^k f)(0^+).$$

Lemma 2.3 [14] Let $\alpha > 0$, If $D_{q,0^+}^\alpha u \in C[0,1]$, then

$$I_{q,0^+}^\alpha D_{q,0^+}^\alpha f(t) = f(t) + \sum_{i=1}^n k_i t^{(\alpha-i)}$$

where $n = [\alpha] + 1$.

Theorem 2.4 (Avery-Henderson fixed point theorem [15]) Let $(E, \|\cdot\|)$ be a Banach space, and $P \subset E$ be a cone. Let Ψ and ϕ be increasing non-negative, continuous functionals on P , and ω be a non-negative

continuous functionals on P with $\omega(0) = 0$, such that, for some $r_3 > 0$ and $M > 0$, $\varphi(u) < \omega(u) < \psi(u)$, and $\|u\| \leq M\varphi(u)$, for all $u \in \overline{p(\varphi, r_3)}$, where $p(\varphi, r_3) = \{u \in P : \varphi(u) < r_3\}$. Suppose that there exist positive numbers $r_1 < r_2 < r_3$, such that

$$\omega(lu) \leq l\omega(u) \quad \text{for } 0 \leq l \leq 1, \text{ and } u \in \partial P(\omega, r_2).$$

If $T : \overline{p(\varphi, r_3)} \rightarrow P$ is a completely continuous operator satisfying:

$$(C_1) \quad \varphi(Tu) > r_3 \text{ for all } u \in \partial P(\omega, r_3);$$

$$(C_2) \quad \varphi(Tu) < r_2 \text{ for all } u \in \partial P(\omega, r_2);$$

$$(C_3) \quad P(\psi, r_1) \neq \emptyset, \text{ and } \psi(Tu) > r_1 \text{ for all } u \in \partial P(\psi, r_1),$$

then T has at least two fixed points u_1 and u_2 such that $r_1 < \psi(u_1)$ with $\omega(u_1) < r_2$ and $r_2 < \omega(u_2)$ with $\varphi(u_2) < r_3$.

III. PRELIMINARY LEMMAS

Lemma 3.1 The boundary value problem (1.1) is equivalent to the following equation:

$$u(t) = d_0 + d_1 t + \frac{1}{\Gamma_q(\alpha)} \int_0^t (t - qs)^{(\alpha-1)} v(s) d_q s, \tag{3.1}$$

where

$$d_0 = \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t - qs)^{(\alpha-1)} v(s) d_q s d_q t + \frac{[b - \int_0^1 t g(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha - 1)} \int_0^1 (1 - qs)^{(\alpha-2)} v(s) d_q s, \tag{3.2}$$

$$d_1 = -\frac{1}{\Gamma_q(\alpha - 1)} \int_0^1 (1 - qs)^{(\alpha-2)} v(s) d_q s, \tag{3.3}$$

$$v(s) = \phi_q \left(\int_0^1 H(s, q\tau) f(\tau, u(\tau)) d_q \tau \right), \tag{3.4}$$

$$H(s, \tau) = \frac{1}{\Gamma_q(\beta)} \begin{cases} (s(1-\tau))^{(\beta-1)} - (s-\tau)^{(\beta-1)}, & 0 \leq \tau \leq s \leq 1, \\ (s(1-\tau))^{(\beta-1)}, & 0 \leq s \leq \tau \leq 1. \end{cases} \tag{3.5}$$

$\phi_q(s)$ is the inverse function of $\phi_p(s)$, a.e., $\phi_q(s) = |s|^{q-2} s$, $\frac{1}{p} + \frac{1}{q} = 1$.

Proof From $D_{q,0^+}^\beta [\phi_p({}^c D_{q,0^+}^\alpha u(t))] + f(t, u(t)) = 0$, we get

$$\phi_p({}^c D_{q,0^+}^\alpha u(t)) = -\frac{1}{\Gamma_q(\beta)} \int_0^t (t - q\tau)^{(\beta-1)} f(\tau, u(\tau)) d_q \tau + c_1 t^{\beta-1} + c_2 t^{\beta-2} + c_3 t^{\beta-3}.$$

In view of $\phi_p({}^c D_{q,0^+}^\alpha u(0)) = [\phi_p({}^c D_{q,0^+}^\alpha u(0))]^\dagger = 0$, we get $c_2 = c_3 = 0$, i.e.,

$$\phi_p({}^c D_{q,0^+}^\alpha u(t)) = -\frac{1}{\Gamma_q(\beta)} \int_0^t (t - q\tau)^{(\beta-1)} f(\tau, u(\tau)) d_q \tau + c_1 t^{\beta-1}. \quad (3.6)$$

Conditions $\phi_p({}^c D_{q,0^+}^\alpha u(1)) = 0$ imply that

$$c_1 = -\frac{1}{\Gamma_q(\beta)} \int_0^1 (1 - q\tau)^{(\beta-1)} f(\tau, u(\tau)) d_q \tau. \quad (3.7)$$

By use of (3.6) and (3.7), we get

$$\phi_p({}^c D_{q,0^+}^\alpha u(t)) = \int_0^1 H(t, q\tau) f(\tau, u(\tau)) d_q \tau. \quad (3.8)$$

In view of (3.8), we obtain

$${}^c D_{q,0^+}^\alpha u(t) = \phi_q \left(\int_0^1 H(t, q\tau) f(\tau, u(\tau)) d_q \tau \right). \quad (3.9)$$

Let

$$v(t) = \phi_q \left(\int_0^1 H(t, q\tau) f(\tau, u(\tau)) d_q \tau \right),$$

by use of (3.9), we get

$$u(t) = \frac{1}{\Gamma_q(\alpha)} \int_0^t (t - qs)^{(\alpha-1)} v(s) d_q s + d_0 + d_1 t + d_2 t^2.$$

Conditions $D_{q,0^+}^2 u(0) = 0$ imply that $d_2 = 0$, i.e.,

$$u(t) = \frac{1}{\Gamma_q(\alpha)} \int_0^t (t - qs)^{(\alpha-1)} v(s) d_q s + d_0 + d_1 t,$$

then we have

$$u'(t) = \frac{1}{\Gamma_q(\alpha-1)} \int_0^t (t - qs)^{(\alpha-2)} v(s) d_q s + d_1.$$

Conditions $D_{q,0^+} u(1) = 0$ imply that

$$d_1 = -\frac{1}{\Gamma_q(\alpha-1)} \int_0^1 (1 - qs)^{(\alpha-2)} v(s) d_q s.$$

From $au(0) + bD_{q,0^+} u(0) = \int_0^1 g(t)u(t) d_q t$, we get

$$\begin{aligned} d_0 &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t - qs)^{(\alpha-1)} v(s) d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 t g(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1 - qs)^{(\alpha-2)} v(s) d_q s. \end{aligned}$$

Therefore, we can obtain

$$\begin{aligned} u(t) &= d_0 + d_1 t + \frac{1}{\Gamma_q(\alpha)} \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s \\ &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 tg(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \\ &\quad - \frac{t}{\Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s + \frac{1}{\Gamma_q(\alpha)} \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s. \end{aligned}$$

The proof is complete. □

Lemma 3.2 ([16]) *The function $H(s, \tau)$ defined by (3.5) is continuous on $[0,1] \times [0,1]$ and satisfy*

$$\frac{s^{\beta-1}(1-s)\tau(1-\tau)^{(\beta-1)}}{\Gamma_q(\beta)} \leq H(s, \tau) \leq \frac{\tau(1-\tau)^{(\beta-1)}}{\Gamma_q(\beta-1)} \quad \text{for } s, \tau \in [0,1].$$

Let E be the real Banach space $C[0,1]$ with the maximum norm, define the operator $T : E \rightarrow E$ by

$$\begin{aligned} Tu(t) &= d_0 + d_1 t + \frac{1}{\Gamma_q(\alpha)} \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s \\ &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 tg(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \\ &\quad - \frac{t}{\Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s + \frac{1}{\Gamma_q(\alpha)} \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s. \end{aligned}$$

Lemma 3.3 *For $u \in C[0,1]$ with $u(t) \geq 0$, $(Tu)(t)$ is non-increasing and non-negative.*

Proof Since

$$Tu(t) = d_0 + d_1 t + \frac{1}{\Gamma_q(\alpha)} \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s,$$

so we get

$$\begin{aligned} (Tu)'(t) &= d_1 + \frac{1}{\Gamma_q(\alpha-1)} \int_0^t (t-qs)^{(\alpha-2)} v(s) d_q s \\ &= -\frac{1}{\Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s + \frac{1}{\Gamma_q(\alpha-1)} \int_0^t (t-qs)^{(\alpha-2)} v(s) d_q s \\ &\leq 0. \end{aligned}$$

So $(Tu)(t)$ is non-increasing, then we have $\min_{t \in [0,1]} Tu(t) = Tu(1)$. We have

$$Tu(1) = d_0 + d_1 + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1-qs)^{(\alpha-1)} v(s) d_q s$$

$$\begin{aligned}
 &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t - qs)^{(\alpha-1)} v(s) d_q s d_q t \\
 &\quad + \frac{[b - \int_0^1 tg(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1 - qs)^{(\alpha-2)} v(s) d_q s \\
 &\quad - \frac{1}{\Gamma_q(\alpha-1)} \int_0^1 (1 - qs)^{(\alpha-2)} v(s) d_q s + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1 - qs)^{(\alpha-1)} v(s) d_q s \\
 &\geq \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t - qs)^{(\alpha-1)} v(s) d_q s d_q t \\
 &\quad + \frac{[b - \int_0^1 tg(t) d_q t] - [a - \int_0^1 g(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1 - qs)^{(\alpha-2)} v(s) d_q s \\
 &\quad + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1 - qs)^{(\alpha-1)} v(s) d_q s \\
 &\geq 0.
 \end{aligned}$$

The proof is complete. □

IV. MAIN RESULTS

Theorem 4.1 Suppose that there exist numbers $0 \leq r_1 \leq r_2 \leq r_3$ such that f satisfies the following conditions:

$$(H_1) \quad f(t, u) > M_3, \quad \text{for } t \in [0, 1], \quad u \in [r_3, \frac{r_3}{k}];$$

$$(H_2) \quad f(t, u) < M_2, \quad \text{for } t \in [0, 1], \quad u \in [0, r_2];$$

$$(H_3) \quad f(t, u) > M_1, \quad \text{for } t \in [0, 1], \quad u \in [0, r_1],$$

where

$$M_3 = \frac{\Gamma_q(\beta)}{qB_q(2, \beta)} \left(\frac{r_3}{L_3} \right)^{\beta-1}, \quad M_2 = \frac{\Gamma_q(\beta-1)}{qB_q(2, \beta)} \left(\frac{r_2}{L_2} \right)^{\beta-1}, \quad M_1 = \frac{\Gamma_q(\beta)}{qB_q(2, \beta)} \left(\frac{r_1}{L_1} \right)^{\beta-1},$$

$$L_3 = \left\{ \frac{b-a}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} + \frac{1}{\Gamma_q(\alpha+1)} \right\} \int_0^1 (s^{\beta-1} (1-s))^{q-1} d_q s,$$

$$L_2 = \frac{\int_0^1 g(t) d_q t}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha+1)} + \frac{b - \int_0^1 tg(t) d_q t}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)},$$

$$L_1 = \frac{b - \int_0^1 tg(t) d_q t}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 (s^{\beta-1} (1-s))^{q-1} d_q s.$$

Then the problem (1.1) has at least two positive solutions u_1 and u_2 such that $r_1 < \psi(u_1)$ with

$\omega(u_1) < r_2$ and $r_2 < \omega(u_2)$ with $\varphi(u_2) < r_3$.

Proof Define the cone $P \subset E$ by

$$P = \left\{ u \mid u \in E, \min_{t \in [0,1]} u(t) \geq k \|u\|, t \in [0,1] \right\},$$

where

$$k = \frac{[b - \int_0^1 t g(t) d_q t] - [a - \int_0^1 g(t) d_q t]}{b - \int_0^1 t g(t) d_q t}, \quad 0 < k < 1.$$

For any $u \in P$, in view of Lemma 3.3, we get

$$\begin{aligned} \min_{t \in [0,1]} |Tu(t)| &= |Tu(1)| = d_0 + d_1 + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1-qs)^{(\alpha-1)} v(s) d_q s \\ &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 t g(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \\ &\quad - \frac{1}{\Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1-qs)^{(\alpha-1)} v(s) d_q s \\ &\geq \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 t g(t) d_q t] - [a - \int_0^1 g(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \\ &\geq k \left\{ \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \right. \\ &\quad \left. + \frac{[b - \int_0^1 t g(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \right\} \\ &= kTu(0) = k \|Tu\|. \end{aligned}$$

Therefore, $T : P \rightarrow P$. In view of the Arzela-Ascoli theorem, we have $T : P \rightarrow P$ is completely continuous.

We define the functions on the cone P :

$$\varphi(u) = \min_{t \in [0,1]} |u(t)| = u(1), \quad \omega(u) = \max_{t \in [0,1]} |u(t)| = u(0),$$

$$\psi(u) = \max_{t \in [0,1]} |u(t)| = u(0).$$

Obviously, we have $\omega(0) = 0$, $\varphi(u) \leq \omega(u) \leq \psi(u)$.

For any $u \in \overline{p(\varphi, r_3)}$, we get $\min_{t \in [0,1]} u(t) \geq k \|u\|$, that is, $\varphi(u) \geq k \|u\|$, therefore we obtain $\|u\| \leq \frac{1}{k} \varphi(u)$.

For any $u \in \partial P(\omega, r_2)$, we get $\omega(lu) = l\omega(u)$ for $0 \leq l \leq 1$.

In the following, we prove that the conditions of Theorem 2.1 hold.

Firstly, let $u \in \partial P(\omega, r_3)$, that is, $u \in [r_3, \frac{r_3}{k}]$ for $t \in [0,1]$. By means of (H_1) , we have

$$\begin{aligned} v(s) &= \phi_q \left(\int_0^1 H(s, q\tau) f(\tau, u(\tau)) d_q \tau \right) \\ &> \phi_q \left(M_3 \int_0^1 \frac{s^{\beta-1} (1-s) q \tau (1-q\tau)^{(\beta-1)}}{\Gamma_q(\beta)} d_q \tau \right) \\ &= \left(\frac{M_3 s^{\beta-1} (1-s) q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q-1}, \end{aligned}$$

where $B_q(2, \beta) = \int_0^1 \tau (1-q\tau)^{(\beta-1)} d_q \tau$. So we get

$$\begin{aligned} \varphi(Tu) &= \min_{t \in [0,1]} |Tu(t)| = Tu(1) = d_0 + d_1 + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1-qs)^{(\alpha-1)} v(s) d_q s \\ &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 tg(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \\ &\quad - \frac{1}{\Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1-qs)^{(\alpha-1)} v(s) d_q s \\ &\geq \frac{[b - \int_0^1 tg(t) d_q t] - [a - \int_0^1 g(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \\ &\quad + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1-qs)^{(\alpha-1)} v(s) d_q s \\ &> \frac{b-a}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} \left(\frac{M_3 s^{\beta-1} (1-s) q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q-1} d_q s \\ &\quad + \frac{1}{\Gamma_q(\alpha)} \int_0^1 (1-qs)^{(\alpha-1)} \left(\frac{M_3 s^{\beta-1} (1-s) q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q-1} d_q s \\ &= \frac{b-a}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \left(\frac{M_3 q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q-1} \int_0^1 (s^{\beta-1} (1-s))^{q-1} d_q s \\ &\quad + \frac{1}{\Gamma_q(\alpha+1)} \left(\frac{M_3 q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q-1} \int_0^1 (s^{\beta-1} (1-s))^{q-1} d_q s \\ &= \left(\frac{M_3 q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q-1} L_3 = r_3. \end{aligned}$$

Secondly, let $u \in \partial P(\omega, r_2)$, that is, $u \in [0, r_2]$ for $t \in [0, 1]$. By means of (H_2) , we get

$$\begin{aligned} v(s) &= \phi_q \left(\int_0^1 H(s, q\tau) f(\tau, u(\tau)) d_q \tau \right) \\ &< \phi_q \left(M_2 \int_0^1 \frac{q\tau(1-q\tau)^{(\beta-1)}}{\Gamma_q(\beta-1)} d_q \tau \right) = \left(\frac{M_2 q B_q(2, \beta)}{\Gamma_q(\beta-1)} \right)^{q-1}. \end{aligned}$$

So we have

$$\begin{aligned} \omega(Tu) &= \max_{t \in [0,1]} |Tu(t)| = Tu(0) = d_0 \\ &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 tg(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} v(s) d_q s \\ &< \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} \left(\frac{M_2 q B_q(2, \beta)}{\Gamma_q(\beta-1)} \right)^{q-1} d_q s d_q t \\ &\quad + \frac{[b - \int_0^1 tg(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha-1)} \int_0^1 (1-qs)^{(\alpha-2)} \left(\frac{M_2 q B_q(2, \beta)}{\Gamma_q(\beta-1)} \right)^{q-1} d_q s \\ &< \frac{\int_0^1 g(t) d_q t}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha+1)} \left(\frac{M_2 q B_q(2, \beta)}{\Gamma_q(\beta-1)} \right)^{q-1} \\ &\quad + \frac{[b - \int_0^1 tg(t) d_q t]}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \left(\frac{M_2 q B_q(2, \beta)}{\Gamma_q(\beta-1)} \right)^{q-1} \\ &= \left(\frac{M_2 q B_q(2, \beta)}{\Gamma_q(\beta-1)} \right)^{q-1} L_2 = r_2. \end{aligned}$$

Finally, let $u \in \partial P(\psi, r_1)$, that is, $u \in [0, r_1]$ for $t \in [0, 1]$. By means of (H_3) , we get

$$\begin{aligned} v(s) &= \phi_q \left(\int_0^1 H(s, q\tau) f(\tau, u(\tau)) d_q \tau \right) \\ &> \phi_q \left(M_1 \int_0^1 \frac{s^{\beta-1} (1-s) q\tau (1-q\tau)^{(\beta-1)}}{\Gamma_q(\beta)} d_q \tau \right) = \left(\frac{M_1 s^{\beta-1} (1-s) q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q-1}. \end{aligned}$$

So we get

$$\begin{aligned} \psi(Tu) &= \max_{t \in [0,1]} |Tu(t)| = Tu(0) = d_0 \\ &= \frac{1}{[a - \int_0^1 g(t) d_q t] \Gamma_q(\alpha)} \int_0^1 g(t) \int_0^t (t-qs)^{(\alpha-1)} v(s) d_q s d_q t \end{aligned}$$

$$\begin{aligned}
 & + \frac{[b - \int_0^1 tg(t)d_q t]}{[a - \int_0^1 g(t)d_q t]\Gamma_q(\alpha - 1)} \int_0^1 (1 - qs)^{(\alpha - 2)} v(s) d_q s \\
 & \geq \frac{[b - \int_0^1 tg(t)d_q t]}{[a - \int_0^1 g(t)d_q t]\Gamma_q(\alpha - 1)} \int_0^1 (1 - qs)^{(\alpha - 2)} v(s) d_q s \\
 & > \frac{[b - \int_0^1 tg(t)d_q t]}{[a - \int_0^1 g(t)d_q t]\Gamma_q(\alpha - 1)} \int_0^1 (1 - qs)^{(\alpha - 2)} \left(\frac{M_1 s^{\beta - 1} (1 - s) q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q - 1} d_q s \\
 & = \frac{[b - \int_0^1 tg(t)d_q t]}{[a - \int_0^1 g(t)d_q t]\Gamma_q(\alpha)} \left(\frac{M_1 q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q - 1} \int_0^1 (s^{\beta - 1} (1 - s))^{q - 1} d_q s \\
 & = \left(\frac{M_1 q B_q(2, \beta)}{\Gamma_q(\beta)} \right)^{q - 1} L_1 = r_1.
 \end{aligned}$$

Therefore, in view of Theorem 2.1, we see that the problem (1.1) has at least two positive solutions u_1 and u_2

such that $r_1 < \psi(u_1)$ with $\omega(u_1) < r_2$ and $r_2 < \omega(u_2)$ with $\varphi(u_2) < r_3$. □

VI. EXAMPLE

In this section, we give a simple example to explain the main result.

Example 5.1 For the problem (1.1), let $\alpha = 2.8$, $\beta = 2.3$, $a = 4$, $b = 10$, $p = 2$, $g(t) = t$, then we get $q = 2$,

$$\int_0^1 g(t)d_q t = \frac{1}{2}, \quad \int_0^1 tg(t)d_q t = \frac{1}{3},$$

$$k = \frac{[b - \int_0^1 tg(t)d_q t] - [a - \int_0^1 g(t)d_q t]}{b - \int_0^1 tg(t)d_q t} = \frac{37}{58} \approx 0.637931.$$

Let

$$f(t, u) = \begin{cases} 23, & t \in [0, 1], u \in [0, 9], \\ 23 + 600(u - 9), & t \in [0, 1], u \in [9, 10], \\ 623, & t \in [0, 1], u \in [10, +\infty). \end{cases}$$

From a direct calculation, we get

$$\begin{aligned}
 f(t, u) & > M_3 \approx 583.266938 & \text{for } & t \in [0, 1], u \in [10, \frac{580}{37}], \\
 f(t, u) & < M_2 \approx 36.538326 & \text{for } & t \in [0, 1], u \in [0, 9], \\
 f(t, u) & > M_1 \approx 21.322041 & \text{for } & t \in [0, 1], u \in [0, 0.5].
 \end{aligned}$$

In view of Theorem 4.1, we see that the aforementioned problem has at least two positive solutions u_1 and u_2

such that $0.5 < \psi(u_1)$ with $\omega(u_1) < 9$ and $9 < \omega(u_2)$ with $\varphi(u_2) < 10$.

REFERENCES

- [1] Chen, T, Liu, W: An anti-periodic boundary value problem for the fractional differential equation with a p -Laplacian operator. *Appl. Math. Lett.* 25, 1671-1675 (2012)
- [2] Chai, G: Positive solutions for boundary value problem of fractional differential equation with p -Laplacian operator. *Bound. Value Probl.* 2012, 18 (2012)
- [3] Feng, X, Feng, H, Tan, H: Existence and iteration of positive solutions for third-order Sturm-Liouville boundary value problems with p -Laplacian. *Appl. Math. Comput.* 266, 634-641 (2015)
- [4] Li, Y, Qi, A: Positive solutions for multi-point boundary value problems of fractional differential equations with p -Laplacian. *Math. Methods Appl. Sci.* 39, 1425-1434 (2015)
- [5] Han, Z, Lu, H, Zhang, C: Positive solutions for eigenvalue problems of fractional differential equation with generalized p -Laplacian. *Appl. Math. Comput.* 257,526-536 (2015)
- [6] Jankowski, T: Positive solutions to fractional differential equations involving Stieltjes integral conditions. *Appl. Math. Comput.* 241, 200-213 (2014)
- [7] Ntouyas, S, Etemad, S: On the existence of solutions for fractional differential inclusions with sum and integral boundary conditions. *Appl. Math. Comput.* 266, 235-243 (2015)
- [8] Cabada, A, Hamdi, Z: Nonlinear fractional differential equations with integral boundary value conditions. *Appl. Math. Comput.* 228, 251-257 (2014)
- [9] Gunendi, M, Yaslan, I: Positive solutions of higher-order nonlinear multi-point fractional equations with integral boundary conditions. *Fract. Calc. Appl. Anal.* 19, 989-1009 (2016)
- [10] Cabada, A, Dimitrijevic, S, Tomovic, T, Aleksic, S: The existence of a positive solution for nonlinear fractional differential equations with integral boundary value conditions. *Math. Methods Appl. Sci.* (2016)
- [11] Yang, W: Positive solution for fractional q -difference boundary value problems with φ -Laplacian operator. *Bull.Malays.Math.Soc.* 36(4)1195-1203(2013)
- [12] Yang, Q, Yang, W: Positive solution for q -fractional four-point boundary value problems with p -Laplacian operator. *J.Inequal.Appl.* 481(2014)
- [13] Bashir, A, Sotiris, K, N, Loannis, K, P: Existence results for nonlocal boundary value problems of nonlinear fractional q -difference equations. *Adv.Differ.Equ.* 140(2012)
- [14] Rajkovic, PM, Marinkovic, SD, Stankovic, MS: Fractional integrals and derivatives in q -calulus. *Appl. Anal. DiscreteMath.* 1,311-323 (2007)
- [15] Avery, RI, Chyan, CJ, Henderson, J: Twin solutions of boundary value problems for ordinary differential equations and finite difference equations. *Comput. Math. Appl.* 42,695-704 (2001)
- [16] Yuan, C: Multiple Positive solutions for $(n-1,1)$ -type semipositone conjugate boundary value problems of nonlinear fractional differential equations. *Electron. J. Qual. Theory Differ. Equ.* 2010, 36 (2010)

Chengmin Hou. "Existence of Positive Solutions for Fractional q -Difference Equations Involving Integral Boundary Conditions with p -Laplacian Operator." *Invention Journal of Research Technology in Engineering & Management (IJRTEM)*, vol. 2, no. 7, 5 July 2018, pp. 05–15., www.ijrtem.com.